

Electromagnetic isoscalar $\rho\pi\gamma$ exchange current and the anomalous action

E. Truhlík^a, J. Smejkal^a and F.C. Khanna^b

^a*Institute of Nuclear Physics, Academy of Sciences of the Czech Republic,
CZ-25068 Řež, Czechia*¹

^b*Theoretical Physics Institute, Department of Physics, University of Alberta,
Edmonton, Alberta, Canada, T6G 2J1 and TRIUMF, 4004 Wesbrook Mall,
Vancouver, BC, Canada, V6T 2A3*²

Abstract

Using modern data, we first refit constants needed to complete an anomalous $\pi\rho\omega a_1$ Lagrangian obtained within the approach of hidden local symmetries. Then we derive from this Lagrangian electromagnetic isoscalar $\rho\pi\gamma$ and $\rho a_1\gamma$ exchange currents needed in calculations of the deuteron electromagnetic form factors at large momentum transfers.

Key words: electromagnetic, isoscalar, exchange, current, anomalous, chiral, Lagrangian

PACS: 11.40.-q, 12.39.Fe, 25.30.Bf

1 Introduction

In calculations of the electromagnetic form factors of the deuteron, the $\rho\pi\gamma$ exchange current plays an important role [1–3]. According to [4–6], it is derived from a $\rho\pi\gamma$ vertex

$$\langle \pi^m(q_2) | J_{\lambda}^{e.m.} | \rho_{\mu}^l(q_1) \rangle = i \frac{e g_{\rho\pi\gamma} K_{\rho\pi\gamma}(q^2)}{m_{\rho}} \varepsilon_{\lambda\nu\sigma\mu} q_{2\nu} q_{\sigma} \delta_{ml}, \quad (1)$$

where $q = q_2 - q_1$ and the constant $g_{\rho\pi\gamma}$ is extracted from the data on the width of the decay $\rho \rightarrow \pi + \gamma$.

¹ E-mail: truhlik@ujf.cas.cz, smejkal@ujf.cas.cz

² E-mail: khanna@phys.ualberta.ca

Here we consider the task of constructing this current by using an anomalous Lagrangian. Such a Lagrangian of the $\pi\rho\omega a_1 f_1$ system was constructed in [7] within the approach of hidden local symmetries. It includes also the external electromagnetic field. The subsystem $\pi\rho\omega a_1$ was considered later in [8], with both the external electromagnetic and weak vector and axial-vector fields included, however. In the Lagrangian, several terms are present, which yield, besides the current (1), a correction to it and also a new current $\rho a_1 \gamma$. The Lagrangian is characterized by several parameters, which should be determined from the data on various reactions. Experimental data available 10 years ago allowed Kaiser and Meissner [7] to extract only some of them. The present experimental situation [9] and results of the recent work [10] allowed us to improve the analysis. As a result, all 4 parameters entering the Lagrangian of the system $\pi\rho\omega a_1$ are now available.

We use further the constructed anomalous Lagrangian to derive the electromagnetic isoscalar $\rho\pi\gamma$ and $\rho a_1 \gamma$ exchange currents, which can be employed in calculations of the electromagnetic form factors of the deuteron. They differ from the standard approach by the presence of an additional momentum dependence in the electromagnetic form factor of the $\rho\pi\gamma$ current and by the completely new $\rho a_1 \gamma$ current. We would like to stress that the considered model fully respects chiral invariance and vector dominance and it is consistent with the present experimental knowledge of elementary processes such as radiative decays $\rho \rightarrow \pi\gamma$, $\omega \rightarrow \pi\gamma$, $f_1 \rightarrow \rho\gamma$ and $f_1 \rightarrow \rho\pi\pi$.

In Sect. 2, we introduce the anomalous Lagrangian, extract the needed constants from the data and present the electromagnetic form factors in our model for the vertices $\rho\pi\gamma$ and $\rho a_1 \gamma$. In Sect. 3, we construct the associated electromagnetic isoscalar exchange currents and make the non-relativistic reduction of these currents. The discussion of the results and our conclusions are given in Sect. 4.

2 Anomalous Lagrangian of the $\pi\rho\omega a_1$ system

We derive the $\rho\pi\gamma$ and $\rho a_1 \gamma$ vertices and the associated exchange current from an anomalous $\pi\rho\omega a_1$ Lagrangian [7,8]. Besides the meson fields, the external vector isoscalar $\mathcal{B}_\mu$ and isovector $\vec{\mathcal{V}}_\mu$ and axial vector isovector $\vec{\mathcal{A}}_\mu$ fields are included. The Lagrangian, which is of interest for the problem under investigation, reads

$$\bar{\mathcal{L}}_7 = 2ig\varepsilon_{\kappa\lambda\mu\nu} \{ \partial_\kappa \omega_\lambda [(g\vec{\rho}_\mu - e\vec{\mathcal{V}}_\mu) \cdot (\frac{1}{f_\pi} \partial_\nu \vec{\pi} + e\vec{\mathcal{A}}_\nu)]$$

$$+ (g\omega_\kappa - \frac{1}{3}e\mathcal{B}_\kappa) [(\partial_\lambda \vec{\rho}_\mu) \cdot (\frac{1}{f_\pi} \partial_\nu \vec{\pi} + e\vec{\mathcal{A}}_\nu)] \}, \quad (2)$$

$$\begin{aligned} \bar{\mathcal{L}}_8 = & -2ig\varepsilon_{\kappa\lambda\mu\nu} \{ \partial_\kappa \omega_\lambda [(g\vec{\rho}_\mu - e\vec{\mathcal{V}}_\mu) \cdot (g\vec{a}_\nu + \frac{1}{2f_\pi} \partial_\nu \vec{\pi})] \\ & + (g\omega_\kappa - \frac{1}{3}e\mathcal{B}_\kappa) [(\partial_\lambda \vec{\rho}_\mu) \cdot (g\vec{a}_\nu + \frac{1}{2f_\pi} \partial_\nu \vec{\pi})] \}, \end{aligned} \quad (3)$$

$$\begin{aligned} \bar{\mathcal{L}}_9 = & 2ie\varepsilon_{\kappa\lambda\mu\nu} \{ \frac{1}{3} \partial_\kappa \mathcal{B}_\lambda [(g\vec{\rho}_\mu - e\vec{\mathcal{V}}_\mu) \cdot (\frac{1}{f_\pi} \partial_\nu \vec{\pi} + e\vec{\mathcal{A}}_\nu)] \\ & + (g\omega_\kappa - \frac{1}{3}e\mathcal{B}_\kappa) [(\partial_\lambda \vec{\mathcal{V}}_\mu) (\frac{1}{f_\pi} \partial_\nu \vec{\pi} + e\vec{\mathcal{A}}_\nu)] \}, \end{aligned} \quad (4)$$

$$\begin{aligned} \bar{\mathcal{L}}_{10} = & -2ie\varepsilon_{\kappa\lambda\mu\nu} \{ \frac{1}{3} \partial_\kappa \mathcal{B}_\lambda [(g\vec{\rho}_\mu - e\vec{\mathcal{V}}_\mu) \cdot (g\vec{a}_\nu + \frac{1}{2f_\pi} \partial_\nu \vec{\pi})] \\ & + (g\omega_\kappa - \frac{1}{3}e\mathcal{B}_\kappa) [(\partial_\lambda \vec{\mathcal{V}}_\mu) \cdot (g\vec{a}_\nu + \frac{1}{2f_\pi} \partial_\nu \vec{\pi})] \}. \end{aligned} \quad (5)$$

Here $g \equiv g_\rho$ and e is the elementary charge. It follows from the KSFR relation, $2f_\pi^2 g^2 = m_\rho^2$, that the value of g is ≈ 6 , if $f_\pi = 92$ MeV and $m_\rho = 770$ MeV are used.

The strong vertices $\rho\omega\pi$ and $\rho\omega a_1$ are already known from the appendix A of Ref. [7]. Then Eqs.(2)-(5) provide all other terms arising from the homogenous terms of the anomalous action due to the presence of the external electroweak interactions which change the natural parity.

The total Yukawa-type anomalous Lagrangian of the $\pi\rho\omega a_1$ system is given by the sum

$$\bar{\mathcal{L}}_{an} = \sum_{i=7,10} \bar{c}_i \bar{\mathcal{L}}_i, \quad (6)$$

where the constants, $\bar{c}_i$, according to [8] are

$$\bar{c}_7 = \tilde{c}_7 + \frac{1}{2}\tilde{c}_8, \quad \bar{c}_8 = \tilde{c}_8, \quad \bar{c}_9 = \tilde{c}_9 + \frac{1}{2}\tilde{c}_{10}, \quad \bar{c}_{10} = \tilde{c}_{10}, \quad (7)$$

and the constants $\tilde{c}_i$ are given in [7] as ¹

$$\tilde{c}_7 = 1.32 \times 10^{-2}, \quad \tilde{c}_8 = -2.05 \times 10^{-1}, \quad \tilde{c}_9 = -5.14 \times 10^{-3}, \quad \tilde{c}_{10} = 0. \quad (8)$$

However, new data on several reactions relevant to the analysis aiming to get these constants has recently been published [9] and also a new work [10], in addition to [7], has appeared, which enables us to get more reliable values of

¹ The constants $\tilde{c}_i$ used here differ from those of Ref. [7] by the factor g .

them. But we still cannot remove an uncertainty in the sign of the constants, implicitly present also in the previous studies [7,10].

2.1 Extracting the constants from the data

We start by extracting the constant $\tilde{c}_7$. It enters the effective constant $g_{\rho\omega\pi}$ defined as

$$\mathcal{L}_{\rho\omega\pi} = i \frac{g_{\rho\omega\pi}}{f_\pi} \varepsilon_{\kappa\lambda\beta\nu} \partial_\kappa \omega_\lambda \vec{\rho}_\beta \cdot \partial_\nu \vec{\pi}. \quad (9)$$

The constant $g_{\rho\omega\pi}$ was found [10] to have the value

$$g_{\rho\omega\pi} = 1.2. \quad (10)$$

On the other hand, our model Lagrangian (6) leads to

$$g_{\rho\omega\pi} = 4g^2 \tilde{c}_7, \quad (11)$$

which yields

$$\tilde{c}_7 = 8.64 \times 10^{-3}. \quad (12)$$

This value of $\tilde{c}_7$ is lower by $\sim 30\%$ than its older value given in (8).

Using the Lagrangian (6) we derive the amplitudes for the radiative processes $\rho \rightarrow \pi\gamma$ and $\omega \rightarrow \pi\gamma$ with the constant $\tilde{c}_9$ entering in combination with $\tilde{c}_7$ into the effective decay constants

$$g_{\rho\pi\gamma} = \frac{2gm_\rho}{3f_\pi}(\tilde{c}_7 + \tilde{c}_9), \quad g_{\omega\pi\gamma} = \frac{2gm_\omega}{f_\pi}(\tilde{c}_7 + \tilde{c}_9). \quad (13)$$

These decay constants enter the radiative decay width of a vector meson V decay into a pseudoscalar meson P as [11]

$$\Gamma_{V \rightarrow P\gamma} = \frac{\alpha}{24} g_{VP\gamma}^2 m_V \left[1 - \left(\frac{m_P}{m_V} \right)^2 \right]^3. \quad (14)$$

According to [9], the widths are

$$\Gamma_{\rho^\pm \rightarrow \pi^\pm \gamma} = (67.82 \pm 7.54) \text{ keV}, \quad \Gamma_{\omega \rightarrow \pi^0 \gamma} = (714.9 \pm 42.1) \text{ keV}. \quad (15)$$

From Eqs. (12)–(15) we have

$$\tilde{c}_9^c = 8.64 \times 10^{-3}, \quad \tilde{c}_9^n = 9.55 \times 10^{-3}. \quad (16)$$

The superscript c (n) implies that the corresponding quantity is obtained from the charged (neutral) meson decay. The values $\tilde{c}_9^c$ and $\tilde{c}_9^n$ differ by $\approx 10\%$.

We calculate the weighted value of $\tilde{c}_9^a$ according to the equation

$$a^w = \left(\sum_{i=1,n} a_i w_i \right) / \left(\sum_{i=1,n} w_i \right), \quad w_i = \Gamma_i / \Delta\Gamma_i. \quad (17)$$

Using Eqs. (15) and (16) we obtain

$$\tilde{c}_9^w = 9.23 \times 10^{-3}. \quad (18)$$

In comparison with $\tilde{c}_9$ from Eq. (8), the new value (18) has the opposite sign and it is larger by a factor of ~ 2 .

If we admit negative values for the sum $\tilde{c}_7 + \tilde{c}_9$ then we get

$$\tilde{c}_9^w = -2.65 \times 10^{-2}. \quad (19)$$

With the weighted value (18) we get for the radiative decay widths

$$\Gamma_{\rho^\pm \rightarrow \pi^\pm \gamma} = 72.5 \text{ keV}, \quad \Gamma_{\omega \rightarrow \pi^0 \gamma} = 689.6 \text{ keV}, \quad (20)$$

and for their ratio

$$\Gamma_{\omega \rightarrow \pi^0 \gamma} / \Gamma_{\rho^\pm \rightarrow \pi^\pm \gamma} = 9.5, \quad (21)$$

and for the effective decay constants

$$g_{\rho\pi\gamma} = 0.585, \quad g_{\omega\pi\gamma} = 1.78. \quad (22)$$

For the negative values, the effective couplings have the opposite sign.

We can also use the same procedure with the neutral ρ meson decay, for which the decay width is [9]

$$\Gamma_{\rho^0 \rightarrow \pi^0 \gamma} = (102.5 \pm 25.6) \text{ keV}. \quad (23)$$

The result for the constant $\tilde{c}_9$ from this equation is

$$\tilde{c}_9 = 1.25 \times 10^{-2}. \quad (24)$$

The experimental value of $\Gamma_{\rho^0 \rightarrow \pi^0 \gamma}$ is converging in time to the value of $\Gamma_{\rho^\pm \rightarrow \pi^\pm \gamma}$ [9], but it is still impossible to get both decay widths at the 1σ error level by using the same value of $\tilde{c}_9$, since the error bounds do not overlap. We can use Eq. (17) for the neutral meson decays to get the weighted value of $\tilde{c}_9^w = 1.014 \times 10^{-2}$, but the calculated value of $\Gamma_{\omega \rightarrow \pi^0 \gamma} = 762 \text{ keV}$ is already outside the 1σ error bound 757 keV . It can be verified that only the values of $\tilde{c}_9$ from the interval $0.97 \times 10^{-2} < \tilde{c}_9 < 1.05 \times 10^{-2}$ yield simultaneously acceptable values of $\Gamma_{\omega \rightarrow \pi^0 \gamma}$ and $\Gamma_{\rho^0 \rightarrow \pi^0 \gamma}$. For

$$\tilde{c}_9 = 1.00 \times 10^{-2}, \quad (25)$$

one gets

$$\Gamma_{\omega \rightarrow \pi^0 \gamma} = 751 \text{ keV}, \quad \Gamma_{\rho^0 \rightarrow \pi^0 \gamma} = 79.4 \text{ keV}, \quad (26)$$

with the ratio

$$\Gamma_{\omega \rightarrow \pi^0 \gamma} / \Gamma_{\rho^0 \rightarrow \pi^0 \gamma} = 9.5, \quad (27)$$

and for the effective decay constants a value of

$$g_{\rho\pi\gamma} = 0.610, \quad g_{\omega\pi\gamma} = 1.86. \quad (28)$$

One obtains similar results with the negative value of $\tilde{c}_9$

$$\tilde{c}_9 = -2.73 \times 10^{-2}. \quad (29)$$

Actually, the decay widths $\Gamma_{\omega \rightarrow \pi^0 \gamma}$ and $\Gamma_{\rho^0 \rightarrow \pi^0 \gamma}$ were used in [10] to extract the constant d , related to our constants by the equation $d = g(\tilde{c}_7 + \tilde{c}_9)$, with the result $d \approx 0.01$. On the other hand, taking our constants g , $\tilde{c}_7$ (12) and $\tilde{c}_9$ (25) we get $d \approx 0.11$. Evidently, the calculations [10] give the value of d^2 instead of d .

It is interesting to follow the change of the value of the constant $g_{\omega\pi\gamma}$ in time. The first use of the $\rho\pi\gamma$ exchange current in the calculations of the electromagnetic deuteron form factors and an estimate of the constant $g_{\omega\pi\gamma}$ was made by Chemtob, Moniz and Rho (CMR) in [5]. Actually, CMR adopted the $\rho\pi\gamma$ exchange current constructed by Adler [4]. The existing data on $\Gamma_{\rho \rightarrow \pi\gamma}$

[12] allowed CMR to extract the upper bound $|g_{\rho\pi\gamma}| < 1.02$. In spite of an estimate, based on the existing relativistic quark model [13], that the value of $g_{\rho\pi\gamma}$ might be about 50% of the upper bound, CMR adopted the value $g_{\rho\pi\gamma} = 1$. Afterwards, the value $g_{\rho\pi\gamma} = 0.34\text{--}0.48$ was extracted by Gari and Hyuga [6] from the more precise data [14]. Later on, Towner [15] analysed the data [16] to obtain the value $g_{\rho\pi\gamma} = 0.578$, which is larger by about 50%. At last, in [18,19], the value of $g_{\rho\pi\gamma}$, obtained from the data [20], has oscillated down to 0.56. However, our value (22) of $g_{\rho\pi\gamma} = 0.585$ is again larger and close to the one obtained in [15]. Towner also extracted from the data [17] the constant $g_{\omega\pi\gamma} = 1.98$, which is larger by about 10% than our $g_{\omega\pi\gamma} = 1.78$ (22).

Let us note that our analysis is based on the data recommended recently in [9] and that in our approach, we extract $g_{\rho\pi\gamma}$ and $g_{\omega\pi\gamma}$ from the data simultaneously. This is due to the fact that in our model, both vertices depend on the same parameters $\tilde{c}_7$ and $\tilde{c}_9$.

We conclude that the present experimental results allow us to extract two values of the constant $\tilde{c}_9$ which differ by $\approx 10\%$ ². However, the difference in the effective coupling constants $g_{\rho\pi\gamma}$ and $g_{\omega\pi\gamma}$ given in Eqs. (22) and (28) is only 4%. One can also ignore the data on the radiative decay $\rho^0 \rightarrow \pi^0 + \gamma$ and use only $\tilde{c}_9$ from Eq. (18) and the decay constants (22) with the hope that the experimental study of this decay will improve.

Let us turn to determining the constants $\tilde{c}_8$ and $\tilde{c}_{10}$, needed to calculate the anomalous processes with the a_1 meson. However, the decays of this meson are not well measured and we use the f_1 meson decays to extract them from the data. In contrast to the experimental situation in late eighties, the decay $f_1 \rightarrow \rho\pi\pi$ is now well measured [9] and good data for the radiative decay $f_1 \rightarrow \rho\gamma$ has appeared [9]. According to [7], one of the contact couplings, $\tilde{h}'_{f_1\rho\pi\pi}$, is connected to $\tilde{c}_8$ as

$$\tilde{c}_8 = -\frac{\tilde{h}'_{f_1\rho\pi\pi}}{g^2}. \quad (30)$$

Using Eq. (4.20) of [7] with the new value

$$\Gamma_{f_1 \rightarrow \rho\pi\pi} = (2.808 \pm 0.360) \text{ MeV}, \quad (31)$$

one gets

$$\tilde{h}'_{f_1\rho\pi\pi} = 3.775, \quad (32)$$

² In the case of the negative values, the uncertainty in $\tilde{c}_9$ is about 3%.

and from (30)

$$\tilde{c}_8 = -1.02 \times 10^{-1}. \quad (33)$$

Again, the new value of $\tilde{c}_8$ is lower by a factor of ~ 2 .

In order to calculate the radiative $f_1 \rightarrow \rho\gamma$ decay, we adopt the pertinent part of the anomalous Lagrangian (3.12) of Ref. [7],

$$\Delta\mathcal{L}_{f_1} = \tilde{c}_8\Delta\mathcal{L}_9 + \tilde{c}_{10}\Delta\mathcal{L}_{12}, \quad (34)$$

$$\Delta\mathcal{L}_9 = -2ig^2\varepsilon_{\kappa\lambda\beta\nu}f_{1\kappa}[g(\partial_\lambda\vec{\rho}_\beta) \cdot \vec{\rho}_\nu - (\partial_\lambda\rho^0)(e\tilde{\mathcal{B}}_\nu)], \quad (35)$$

$$\mathcal{L}_{12} = -2ig^2\varepsilon_{\kappa\lambda\beta\nu}f_{1\kappa}(\partial_\lambda e\tilde{\mathcal{B}}_\beta)\rho_\nu^0. \quad (36)$$

With these vertices, we get for the radiative decay width an equation similar to Eq. (14)

$$\Gamma_{f_1 \rightarrow \rho\gamma} = \frac{\alpha}{24}g_{\rho f_1 \gamma}^2 m_{f_1} \left[1 + \left(\frac{m_{f_1}}{m_\rho} \right)^2 \right] \left[1 - \left(\frac{m_\rho}{m_{f_1}} \right)^2 \right]^3, \quad (37)$$

where

$$g_{\rho f_1 \gamma} = 2g^2 (\tilde{c}_8 + \tilde{c}_{10}). \quad (38)$$

Using the experimental value of $\Gamma_{f_1 \rightarrow \rho\gamma}$ [9]

$$\Gamma_{f_1 \rightarrow \rho\gamma} = (1.296 \pm 0.288) \text{ MeV}, \quad (39)$$

we get

$$\tilde{c}_8 + \tilde{c}_{10} = 2.65 \times 10^{-2}, \quad (40)$$

and for the constant $\tilde{c}_{10}$ the value

$$\tilde{c}_{10} = 1.29 \times 10^{-1}, \quad (41)$$

which is in absolute value even larger than $\tilde{c}_8$, Eq. (33). Then for the effective coupling $g_{f_1 \rho \gamma}$ we have the value

$$g_{f_1 \rho \gamma} = 1.84. \quad (42)$$

Admitting negative values for the sum $\tilde{c}_8 + \tilde{c}_{10}$ we get

$$\tilde{c}_{10} = 7.59 \times 10^{-2}. \quad (43)$$

We now have in Eqs. (12),(18),(25),(33) and (41) all the needed constants and we can now proceed to constructing the $\rho\pi\gamma$ and $\rho a_1\gamma$ exchange currents. However, let us first discuss the electromagnetic form factors provided by our model.

2.2 Electromagnetic form factors

The Lagrangian of Eq. (6) yields the vertex $\rho\pi\gamma$, Eq. (1), with the form factor

$$K_{\rho\pi\gamma}(q^2) = F_\omega(q^2) + \left(\frac{\tilde{c}_9 - \tilde{c}_7}{\tilde{c}_9 + \tilde{c}_7} \right) [1 - F_\omega(q^2)], \quad (44)$$

where the form factor $F_\omega(q^2)$ is given by the vector dominance. Since $\tilde{c}_9 > \tilde{c}_7$, the second term on the right hand side of Eq. (44) gives a 3–7% positive correction to the first term at $\vec{q}^2 \approx m_\omega^2$. In the case of the negative values of $\tilde{c}_9$, this term is larger by a factor of ~ 2 than the first one at these values of $\vec{q}^2$.

The presence of the terms $\bar{\mathcal{L}}_8$ and $\bar{\mathcal{L}}_{10}$ in Eq. (6) shows the existence of the vertices $\rho a_1\gamma$, which give rise to the corresponding $\rho a_1\gamma$ currents. In analogy with Eq. (1) we have

$$\langle a_{1\nu}^m(q_2) | J_\lambda^{e.m.} | \rho_\mu^l(q_1) \rangle = e g_{\rho a_1\gamma} \varepsilon_{\lambda\nu\sigma\mu} [K_1(q^2) q_\sigma + K_2(q^2) q_{1\sigma}] \delta_{ml}, \quad (45)$$

where the form factors $K_{1,2}(q^2)$ are defined as

$$K_1(q^2) = \frac{1}{\tilde{c}_8 + \tilde{c}_{10}} [\tilde{c}_8 F_\omega(q^2) + \tilde{c}_{10}], \quad K_2(q^2) = \frac{\tilde{c}_8}{\tilde{c}_8 + \tilde{c}_{10}} [F_\omega(q^2) - 1]. \quad (46)$$

The effective coupling constant $g_{\rho a_1\gamma}$ is

$$g_{\rho a_1\gamma} = g_{\rho f_1\gamma}/3, \quad (47)$$

with the constant $g_{\rho f_1\gamma}$ given in Eq. (38). This gives a new contribution to the current.

In the next section, we derive the $\rho\pi\gamma$ and $\rho a_1\gamma$ exchange currents.

3 The $\rho\pi\gamma$ and $\rho a_1\gamma$ exchange currents

The general structure of the exchange currents which follows from our Lagrangian (6) is given in Fig. 1. The currents of the pion range are,

$$J_{\pi\mu}^{m.c.} = ig_{\pi NN} \frac{2g^2\tilde{c}_7}{3f_\pi} \varepsilon_{\mu\kappa\beta\nu} F_\omega(q^2) q_\kappa \mathcal{P}_\beta(1,2) q_{2\nu} + (1 \leftrightarrow 2), \quad (48)$$

$$J_{\pi\mu}^c = -ig_{\pi NN} \frac{g^2(\tilde{c}_7 - \tilde{c}_9)}{3f_\pi} \varepsilon_{\mu\kappa\beta\nu} q_\kappa \mathcal{P}_\beta(1,2) q_{2\nu} + (1 \leftrightarrow 2), \quad (49)$$

where

$$\mathcal{P}_\beta(1,2) = \Gamma_{1,\beta}^a \Delta_F^\rho(q_1^2) \Delta_F^\pi(q_2^2) \Gamma_{2,5}^a, \quad (50)$$

and

$$\Gamma_{i,\beta}^a = \bar{u}(p'_i)(\gamma_\beta - \frac{\kappa_V}{2M}\sigma_{\beta\delta} q_{i\delta})\tau^a u(p_i) \text{ and } \Gamma_{j,5}^a = \bar{u}(p'_j)\gamma_5\tau^a u(p_j). \quad (51)$$

The sum of the currents given by Eqs. (48) and (49) is

$$J_{\pi\mu} = ig_{\pi NN} \frac{g}{2} \frac{g_{\rho\pi\gamma}}{m_\rho} K_{\rho\pi\gamma}(q^2) \varepsilon_{\mu\kappa\beta\nu} q_\kappa \mathcal{P}_\beta(1,2) q_{2\nu} + (1 \leftrightarrow 2), \quad (52)$$

where the effective coupling constant $g_{\rho\pi\gamma}$ and the form factor $K_{\rho\pi\gamma}(q^2)$ are defined in Eqs. (13) and (44), respectively.




Fig. 1. The general structure of the electromagnetic isoscalar $\rho\pi\gamma$ and $\rho a_1\gamma$ exchange currents $J_\mu(q)$ considered in this paper. The meson B_1 is the ρ meson.

The range of the current is given by the meson B_2 which is either π or a_1 meson. The graph a is for a mesonic current $J_{B_2\mu}^{m.c.}$, where the electromagnetic isoscalar interaction is mediated by the meson B which is ω meson. Contact terms are given by the graph b, $J_{B_2\mu}^c$, where the isoscalar electromagnetic current interacts directly with the mesons B_1 and B_2 .

The currents of the a_1 meson range are

$$J_{a_1\mu}^{m.c.} = -\frac{2g_A}{3} g^4 \tilde{c}_8 \varepsilon_{\mu\kappa\beta\nu} F_\omega(q^2) (q_\kappa + q_{1\kappa}) \mathcal{P}_{\beta\nu}(1, 2) + (1 \leftrightarrow 2), \quad (53)$$

$$J_{a_1\mu}^c = \frac{g_A}{3} g^4 \varepsilon_{\mu\kappa\beta\nu} (\tilde{c}_8 q_{1\kappa} - \tilde{c}_{10} q_\kappa) \mathcal{P}_{\beta\nu}(1, 2) + (1 \leftrightarrow 2), \quad (54)$$

where the weak axial coupling constant $g_A = 1.26$, and

$$\mathcal{P}_{\beta\nu}(1, 2) = \Gamma_{1,\beta}^a \Delta_F^\rho(q_1^2) \Delta_F^{a_1}(q_2^2) \Gamma_{2,5\nu}^a \text{ and } \Gamma_{j,5\nu}^a = \bar{u}(p'_j) \gamma_\nu \gamma_5 \tau^a u(p_j). \quad (55)$$

The total $\rho a_1 \gamma$ current is

$$J_{a_1\mu} = -\frac{g_A}{2} g^2 g_{\rho a_1 \gamma} \varepsilon_{\mu\kappa\beta\nu} \left[K_1(q^2) q_\kappa + K_2(q^2) q_{1\kappa} \right] \mathcal{P}_{\beta\nu}(1, 2) + (1 \leftrightarrow 2), \quad (56)$$

with the effective coupling constant $g_{\rho a_1 \gamma}$ and with the form factors $K_{1,2}(q^2)$ given in Eqs. (47) and (46), respectively.

In deriving Eqs. (48) and (49) and Eqs. (53) and (54), we use the equation $g_\omega = 3g_\rho$, which ensures the current conservation.

3.1 Non-relativistic reduction

Now we proceed to the non-relativistic reduction of the currents given by Eqs. (52) and (56). Having in mind the application of our results to the elastic electron scattering by nuclei, we neglect the energy transfer putting $q_0 = 0$. This reduction yields for the space component of the currents

$$\begin{aligned} \vec{J}_\pi &= C_\pi \left\langle \left(\vec{q} \times \vec{q}_2 \right) - \frac{1}{4M^2} \left\{ \left(\frac{1}{2} + \kappa_V \right) \left(\vec{q} \times \vec{q}_2 \right) \left[\vec{q}_1^2 + i \vec{\sigma}_1 \cdot \vec{P}_1 \times \vec{q}_1 \right] \right. \right. \\ &\quad \left. \left. - i \left(1 + \kappa_V \right) \left(\vec{q}_2 \cdot \vec{P}_2 \right) \left(\vec{q} \times \left(\vec{\sigma}_1 \times \vec{q}_1 \right) \right) \right\} \right\rangle \\ &\quad \cdot \left(\vec{\sigma}_2 \cdot \vec{q}_2 \right) \left(\vec{\tau}_1 \cdot \vec{\tau}_2 \right) \Delta_F^\rho(\vec{q}_1^2) \Delta_F^\pi(\vec{q}_2^2) + (1 \leftrightarrow 2), \\ \vec{J}_{a_1} &= i \frac{g_A}{2} g^2 g_{\rho a_1 \gamma} \left(K_1(\vec{q}^2) \vec{q} + K_2(\vec{q}^2) \vec{q}_1 \right) \times \left\langle \vec{\sigma}_2 \left\{ 1 - \frac{1 + 2\kappa_V}{8M^2} \left[\vec{q}_1^2 \right. \right. \right. \\ &\quad \left. \left. \left. + i \vec{\sigma}_1 \cdot \vec{P}_1 \times \vec{q}_1 \right] \right\} - i \left(\vec{\sigma}_1 \times \vec{q}_1 \right) \frac{1 + \kappa_V}{4M^2} \left(\vec{\sigma}_2 \cdot \vec{P}_2 \right) \right\rangle \end{aligned} \quad (57)$$

$$\cdot (\vec{\tau}_1 \cdot \vec{\tau}_2) \Delta_F^\rho(\vec{q}_1^2) \Delta_F^{a_1}(\vec{q}_2^2) + (1 \leftrightarrow 2). \quad (58)$$

Here

$$C_\pi = -i \frac{g_{\pi NN}}{2M} \frac{g}{2m_\rho} g_{\rho\pi\gamma} K_{\rho\pi\gamma}(\vec{q}^2), \quad (59)$$

and

$$\vec{P}_i = \vec{p}_i + \vec{p}_i. \quad (60)$$

The importance of relativistic corrections in the space component of the $\rho\pi\gamma$ current was stressed in [18]. They were obtained earlier in [21]. We present them also in the a_1 exchange term. An evaluation of the contribution of these terms to the electromagnetic deuteron form factors would give more confidence in the validity of the static approximation for the current (58).

Besides the relativistic corrections present in the current (57), we list the boost current $\vec{J}_\pi^B$ arising from the commutator of the leading terms in Eq. (57) with the operator of the kinematical boost and the retardation current $\vec{J}_\pi^{ret}$. The kinematical boost current is defined as [22]

$$\vec{J}_\pi^B = i [\chi_0(\vec{P}), \vec{J}_\pi^0], \quad (61)$$

where

$$\chi_0(\vec{P}) = -\frac{i}{16M^2} [-\vec{P}^2 - 2i(\vec{r} \cdot \vec{P})(\vec{p} \cdot \vec{P}) + 2i(\vec{\sigma}_1 - \vec{\sigma}_2) \times \vec{p} \cdot \vec{P}], \quad (62)$$

$$\vec{r} = \vec{r}_1 - \vec{r}_2, \quad (63)$$

and the static approximation to the $\rho\pi\gamma$ current $\vec{J}_\pi^0$ is defined as

$$\vec{J}_\pi^0 = C_\pi (\vec{q} \times \vec{q}_2) (\vec{\sigma}_2 \cdot \vec{q}_2) (\vec{\tau}_1 \cdot \vec{\tau}_2) \Delta_F^\rho(\vec{q}_1^2) \Delta_F^\pi(\vec{q}_2^2) + (1 \leftrightarrow 2). \quad (64)$$

Generally, the current $\vec{J}_\pi^B$ is quite complicated. However, it simplifies in the Breit system, where it reads in the c.m. coordinates,

$$\begin{aligned} \vec{J}_\pi^B = & \frac{i}{16M^2} C_\pi (\vec{q} \times \vec{q}_-) \left\{ (\vec{\sigma}_2 \cdot \vec{q}_-) \left[-\frac{1}{4} (\vec{r} \cdot \vec{q}) (\vec{q} \cdot \vec{k}) + (\vec{\sigma}_1 \cdot \vec{K} \times \vec{q}) \right] \right. \\ & - (\vec{q}_- \cdot \vec{K} \times \vec{q}) + i (\vec{\sigma}_2 \times \vec{q}_-) \cdot (\vec{k} \times \vec{q}) \left. \right\} \\ & \cdot (\vec{\tau}_1 \cdot \vec{\tau}_2) \Delta_F^\rho(\vec{q}_+^2) \Delta_F^\pi(\vec{q}_-^2) + (1 \leftrightarrow 2). \end{aligned} \quad (65)$$

Here

$$\begin{aligned}
\vec{p} &= \frac{1}{2}(\vec{p}_1 - \vec{p}_2), \quad \vec{p}' = \frac{1}{2}(\vec{p}'_1 - \vec{p}'_2), \quad \vec{P} = \vec{p}_1 + \vec{p}_2, \quad \vec{P}' = \vec{p}'_1 + \vec{p}'_2, \\
\vec{k} &= \vec{p}' - \vec{p}, \quad \vec{K} = \vec{p}' + \vec{p}, \quad \vec{q} = \vec{P}' - \vec{P}, \quad \vec{Q} = \vec{P}' + \vec{P}, \\
\vec{q}_1 &= \vec{p}'_1 - \vec{p}_1 = \vec{k} + \frac{1}{2}\vec{q} \equiv \vec{q}_+, \quad \vec{q}_2 = \vec{p}'_2 - \vec{p}_2 = -\vec{k} + \frac{1}{2}\vec{q} \equiv -\vec{q}_-, \\
\vec{P}_1 &= \vec{p}'_1 + \vec{p}_1 = \vec{K} + \frac{1}{2}\vec{Q}, \quad \vec{P}_2 = \vec{p}'_2 + \vec{p}_2 = -\vec{K} + \frac{1}{2}\vec{Q}.
\end{aligned} \tag{66}$$

In the Breit system, $\vec{Q} = \vec{P}' - \vec{P} = 0$ and $\vec{P}' = -\vec{P} = \vec{q}/2$.

In the realistic calculations, the currents, Eq. (52) and Eq. (56), are multiplied by the strong form factors. In essence, this procedure reduces in the case of the $\rho\pi\gamma$ current to the following change in Eq. (50),

$$\Delta_F^\rho(\vec{q}_1^2) \Delta_F^\pi(\vec{q}_2^2) \rightarrow F_{\rho NN}(\vec{q}_1^2) \Delta_F^\rho(\vec{q}_1^2) F_{\pi NN}(\vec{q}_2^2) \Delta_F^\pi(\vec{q}_2^2), \tag{67}$$

and to the change

$$\Delta_F^\rho(\vec{q}_1^2) \Delta_F^{a_1}(\vec{q}_2^2) \rightarrow F_{\rho NN}(\vec{q}_1^2) \Delta_F^\rho(\vec{q}_1^2) F_{a_1 NN}(\vec{q}_2^2) \Delta_F^{a_1}(\vec{q}_2^2), \tag{68}$$

in Eq. (55) in the case of the $\rho a_1 \gamma$ current. While in this case, the static approximation is expected to be reasonably good, for the $\rho\pi\gamma$ current the retardation current arises by the following change in the current $\vec{J}_\pi^0$,

$$\begin{aligned}
\Delta_F^\rho(\vec{q}_1^2) \Delta_F^\pi(\vec{q}_2^2) &\rightarrow F_{\rho NN}(\vec{q}_1^2) \Delta_F^\rho(\vec{q}_1^2) F_{\pi NN}(\vec{q}_2^2) \Delta_F^\pi(\vec{q}_2^2) \\
&\times \left\{ 1 + q_{10}^2 \left[\Delta_F^\rho(\vec{q}_1^2) - \frac{\partial}{\partial \vec{q}_1^2} \ln(F_{\rho NN}(\vec{q}_1^2)) \right] \right. \\
&\left. + q_{20}^2 \left[\Delta_F^\pi(\vec{q}_2^2) - \frac{\partial}{\partial \vec{q}_2^2} \ln(F_{\pi NN}(\vec{q}_2^2)) \right] \right\}.
\end{aligned} \tag{69}$$

The retardation current $\vec{J}_\pi^{ret}$ is proportional to the energy transfers q_{10} and q_{20} defined as

$$q_{i0} = \frac{1}{2M} \vec{q}_i \cdot \vec{P}_i, \tag{70}$$

which can be expressed via the c.m. variables by using Eqs. (66). The part of the current $\vec{J}_\pi^{ret}$ without the contribution from the derivative of the form factors was derived earlier in [21].

The exchange charge densities are

$$J_{\pi 0} = \frac{g_{\pi NN}}{2M} \frac{g}{2m_\rho} \frac{1 + \kappa_V}{2M} g_{\rho\pi\gamma} K_{\rho\pi\gamma}(\vec{q}^2) [\vec{q} \cdot \vec{q}_2 \times (\vec{\sigma}_1 \times \vec{q}_1)] \cdot (\vec{\tau}_1 \cdot \vec{\tau}_2) \Delta_F^\rho(\vec{q}_1^2) \Delta_F^\pi(\vec{q}_2^2) + (1 \leftrightarrow 2), \quad (71)$$

$$J_{a_1 0} = \frac{g_A}{2} g^2 g_{\rho a_1 \gamma} \frac{1 + \kappa_V}{2M} \left[K_1(\vec{q}^2) \vec{q} + K_2(\vec{q}^2 \vec{q}_1) \right] \cdot [\vec{\sigma}_2 \times (\vec{q}_1 \times \vec{\sigma}_1)] \cdot (\vec{\tau}_1 \cdot \vec{\tau}_2) \Delta_F^\rho(\vec{q}_1^2) \Delta_F^{a_1}(\vec{q}_2^2) + (1 \leftrightarrow 2). \quad (72)$$

4 Results and conclusions

In this paper, we have studied the electromagnetic processes in the $\pi\rho a_1\omega$ system, characterized by the change of the natural parity of the participating particles, by using the anomalous chiral Lagrangian [7,8]. This Lagrangian contains 4 constants $\tilde{c}_7$ – $\tilde{c}_{10}$, which were first extracted from the data in [7]. Using new data [9] on the radiative decays of the ρ^- , ω^- and f_1 mesons and on the decay $f_1 \rightarrow \rho\pi\pi$ and the constant $\tilde{c}_7$ from [10], we have determined the constants $\tilde{c}_8$, $\tilde{c}_9$ and $\tilde{c}_{10}$. The new constants differ considerably from the old ones. These constants enter the decay rates in the combinations $\tilde{c}_7 + \tilde{c}_9$ and $\tilde{c}_8 + \tilde{c}_{10}$, which can be obtained only up to the sign. We considered both possibilities and we extracted two sets of constants. Disregarding the sign problem, these constants are now known within a 10–15% accuracy. The main source of the uncertainty is due to an inconsistency in the data for the radiative decays of the charged and neutral ρ mesons and of the ω meson. However, the effective couplings $g_{\rho\pi\gamma}$ and $g_{\omega\pi\gamma}$ suffer from this inconsistency by an uncertainty only of about 4%.

We have subsequently constructed the electromagnetic form factor $K_{\rho\pi\gamma}(q^2)$ that contains, besides the well known term, an additional piece which can be large, if the corresponding choice of the constant $\tilde{c}_9$ is made. The presence of the $\rho a_1\omega$ vertex in the Lagrangian gives rise to the new $\rho a_1\gamma$ vertex, which contains two form factors $K_1(q^2)$ and $K_2(q^2)$.

The isoscalar exchange currents, associated with these vertices, are constructed as well. In addition to the known $\rho\pi\gamma$ current, we have also $\rho a_1\gamma$ current with well defined parameters extracted from the data. This is in contrast to $\omega\epsilon\gamma$ [18] or $\omega\sigma\gamma$ [19] currents based on less firm ground. After the non-relativistic reduction, relativistic corrections are also preserved in the space component of our currents, in conformity with the results of Refs. [18],[21]. As to the $\rho\pi\gamma$ current, the leading terms on the right hand sides of Eqs. (71) and (57) are the same as the exchange charge density (3. 2a) and the space component (3.2b) in [6], respectively. However, our electromagnetic form factor $K_{\rho\pi\gamma}(q^2)$ Eq. (44) exhibits an additional momentum dependence. Our relativistic corrections contain the contributions from the vertices, from the kinematical boost and from the retardation, with the part arising from the strong form factors

included. The current $\vec{J}_{a_1}$ Eq. (58) and the density $J_{a_1 0}$ Eq. (72) are entirely new. The subsequent numerical work should reveal how much they can influence the deuteron form factors at high momentum transfers.

Acknowledgments

This work is supported in part by the grant GA ĆR 202/00/1669. A part of this work was done during the stay of E. T. at the Theoretical Physics Institute of the University of Alberta. He thanks Prof. F. C. Khanna for the warm hospitality. The research of F. C. K. is supported in part by NSERCC.

We thank Prof. B. Mosconi for the correspondence and R. Teshima for the discussion.

References

- [1] S. Platchkov *et al.*, *Nucl. Phys. A* **510** (1990) 740.
- [2] B. Mosconi and P. Ricci, *Few-Body Systems* **6** (1989) 63.
- [3] P. Sarriguren, J. Martorell and D. Sprung, *Phys. Lett. B* **228** (1989) 285.
- [4] R.J. Adler, *Phys. Rev.* **141** (1966) 1499.
- [5] M. Chemtob, E. Moniz and M. Rho, *Phys. Rev. C* **10** (1974) 344.
- [6] M. Gari and H. Hyuga, *Nucl. Phys. A* **264** (1976) 409.
- [7] N. Kaiser and U.-G. Meißner, *Nucl. Phys. A* **519** (1990) 671.
- [8] J. Smejkal, E. Truhlík and F. C. Khanna, Non-abelian anomaly and the Lagrangian for radiative muon capture, in: J. Adam, Jr. *et al.*, eds., *Proceedings Sevenths International Conference 'Mesons and Light Nuclei 98', Prague-Průhonice, 1998* (World Scientific, Singapore, 1999) 490.
- [9] C. Caso *et al.*, (Particle Data Group), *Eur. Phys. J. C* **3** (1998) 1.
- [10] F. Klingl, N. Kaiser and W. Weise, *Z. Phys. A* **356** (1996) 193.
- [11] O. Dumbrajs *et al.*, *Nucl. Phys. B* **216** (1983) 277.
- [12] T.A. Lasinski *et al.*, (Particle Data Group), *Rev. Mod. Phys. (Suppl.)* **45** (1973) S1.
- [13] R. Feynman, M. Kislinger and F. Ravndal, *Phys. Rev. D* **3** (1971) 2706.
- [14] B. Gobbi *et al.*, *Phys. Rev. Lett.* **33** (1974) 1450.

- [15] I.S. Towner, *Phys. Rep.* **155** (1987) 263.
- [16] T. Jensen *et al.*, *Phys. Rev. D* **27** (1983) 26.
- [17] M. Aguilar-Benitez *et al.*, (Particle Data Group), *Phys. Lett. B* **170** (1986) 1.
- [18] E. Hummel and J.A. Tjon, *Phys. Rev. C* **42** (1990) 423.
- [19] H. Ito and F. Gross, *Phys. Rev. Lett.* **71** (1993) 2555.
- [20] D. Berg *et al.*, *Phys. Rev. Lett.* **44** (1980) 706.
- [21] K. Tamura, T. Niwa, T. Sato and H. Ohtsubo, *Nucl. Phys. A* **536** (1992) 597.
- [22] J. Adam, Jr. and H. Arenhövel, *Nucl. Phys. A* **614** (1997) 289.